

Planar Binomial Coefficients

Lothar Gerritzen

06.01.2005

Abstract

The notion of binomial coefficients $\binom{T}{S}$ of finite planar, reduced rooted trees T, S is defined and a recursive formula for its computation is shown. The nonassociative binomial formula

$$(1+x)^T = \sum_S \binom{T}{S} x^S$$

for powers relative to T is derived.

Similarly binomial coefficients $\binom{T}{S,V}$ of the second kind are introduced and

it is shown that $(x \otimes 1 + 1 \otimes x)^T = \sum_{S,V} \binom{T}{S,V} (x^S \otimes x^V)$

The roots $\sqrt[T]{1+x} = (1+x)^{T^{-1}}$ which are planar power series f such that $f^T = 1+x$ are considered. Formulas for their coefficients are given.

1 Introduction

The binomial coefficient $\binom{T}{S}$ of two finite, planar, reduced rooted trees T, S is the number of contractions of T onto S . If C_r denotes r -corona, then $\binom{C_n}{C_m} = \binom{n}{m}$. Also

$$\sum_S \binom{T}{S} = \binom{n}{m}$$

if $\deg(T) = n$ and the summation is extended over all S with $\deg(S) = m$ where the degree $\deg(T)$ is the number of leaves of T .

If $(1+x)^T$ denotes the T -th power of $1+x$, then we obtain the nonassociative binomial formula

$(1+x)^T = \sum_S \binom{T}{S} (x^S)$ where the summation is extended over all trees S .

The T -th root $\sqrt[T]{1+x}$ is the unique power series f with constant coefficient 1 and $f^T = 1+x$.

It is uniquely defined and its coefficients are denoted by $\binom{T^{-1}}{S}$. They are computed

in section 6, if T is a corona.

It is an open problem to device a procedure for the computation of the coefficients $\binom{T^{-1}}{S}$ of $\sqrt[3]{1+x}$.

Also we consider binomial coefficients $\binom{T}{S,V}$ of the second kind which are defined to be the number of contractions of T onto S for which the complementary contractions are onto V . It is shown that

$$(x \otimes 1 + a \otimes x)^T = \sum \binom{T}{S,V} x^S \otimes x^V$$

2 Binomial coefficients of the first kind

Denote by $\mathbb{P}$ the set of isomorphism classes of finite, planar, reduced rooted trees. The empty tree $\underline{1}$ is included in $\mathbb{P}$. For $T \in \mathbb{P}$ and any subset I of the set $L(T)$ of leaves of T denote by $T|I$ the contraction of T onto I , see [G 3].

Definition 2.1. For any pair T, S of $\mathbb{P}$ denote by

$$\binom{T}{S}$$

the number of $I \subseteq L(T)$ such that the contraction $T|I$ of T onto I is isomorphic with S .

We call $\binom{T}{S}$ the planar binomial coefficient of T over S of the first kind

Proposition 2.2. The following elementary properties are easy to check:

- (i) $\binom{T}{S} = 0$, if $\deg(S) > \deg(T)$
- (ii) If $\deg(S) = \deg(T)$, then $\binom{T}{S} = 1$ if $S = T$ and $\binom{T}{S} = 0$ if $S \neq T$.
- (iii) $\binom{T}{1} = 1$
- (iv) If x denotes the tree with a single vertex, then $\binom{T}{x} = \deg(T)$
- (v) $\binom{T}{x^2} = \binom{n}{2}$, if $n = \deg(T)$ and if x^2 denotes the unique tree of degree 2 in $\mathbb{P}$.

$$(vi) \quad \sum_{S \in \mathbb{P}, \deg(S)=m} \binom{T}{S} = \deg \binom{(T)}{m}$$

Let $T_1, \dots, T_m \in \mathbb{P}$ and denote by $\cdot_m(T_1, \dots, T_m)$ or $T_1 \cdot T_2 \cdots T_m$ the m -ary grafting of $T_1, \dots, T_m$. It is the unique tree T in $\mathbb{P}$ for which $T - \rho_T$ is the planar forest which is the disjoint union of the ordered systems $T_1 \dot{\bigcup} \dots \dot{\bigcup} T_m$ where T_i is less than T_j in case i is less than j . Here ρ_T is the root of T .

Proposition 2.3. Let $m, r \geq 2$ and $T = T_1 \cdot T_2 \cdot \dots \cdot T_m$ and $S = S_1 \cdot \dots \cdot S_r \in \mathbb{P}$ and $T_i \neq \underline{1}, S_j \neq \underline{1}$ for all i, j .

Then we have the following recursion formula

$$\binom{T}{S} = \sum_{1 \leq i_1 < i_2 < \dots < i_r \leq m} \binom{T_{i_1}}{S_1} \cdot \dots \cdot \binom{T_{i_r}}{S_r} + \sum_{i=1}^m \binom{T_i}{S}$$

Example 2.4. Let $T = T_1 \cdot \dots \cdot T_m, \deg(T_i) = n_i$, then

$$\binom{T}{xx^2} = \sum_{i=1}^m \binom{T_i}{xx^2} + \sum_{1 \leq i < j \leq m} n_i \binom{n_j}{2}$$

$$\binom{T}{x^2x} = \sum_{i=1}^m \binom{T_i}{x^2x} + \sum_{1 \leq i < j \leq m} \binom{n_i}{2} n_j$$

and

$$\binom{n}{3} = \sum_{i=1}^m \binom{n_i}{3} + \sum_{1 \leq i < j \leq m} \frac{n_i n_j}{2} (n_i + n_j - 2)$$

If $m = 2$, then $n_1 + n_2 = n$

and

$$\binom{n}{3} = \binom{n_1}{3} + \binom{n_2}{3} + \frac{n_1 n_2}{2} (n_1 + n_2 - 2)$$

3 Coefficients of the second kind

Let $T, S, V \in \mathbb{P}$.

Definition 3.1. The number

$$\binom{T}{S, V} := \#\{I \subseteq L(T) : T|I = S, T|(L(T) - I) = V\}$$

is called the planar binomial coefficient of T over S and V of the second kind.

In combinatorics there are well-known Stirling numbers of the first and second kind which has been a reason for choosing this terminology.

From the definition it immediately follows that $\sum_{V \in \mathbb{P}} \binom{T}{S, V} V = \binom{T}{S}$

and that $\binom{T}{S, V} = \binom{T}{V, S}$

For $m, r, s \geq 2$ denote by $\Gamma(m, r, s)$ the set of all pairs (α, β) of subsets of $\underline{m} := \{1, 2, \dots, m\}$ with the following properties.

(i) $\#\alpha \in \{1, r\}, \#\beta \in \{1, s\}$

(ii) $\alpha \cup \beta = \{1, 2, \dots, m\}$

Then $\Gamma'(m, r, s) := \{(\alpha, \beta) \in \Gamma(m, r, s) : \#\alpha = 1\}$
is equal to $\{(\{i\}, \underline{m}) : 1 \leq i \leq r\}$ if $s = m$ and is equal to

$$\{(\{i\}, \underline{m} - \{i\}) : 1 \leq i \leq r\}$$

if $s = m - 1$ and $\Gamma'(m, r, s)$ is empty otherwise.

Also $\Gamma''(m, r, s) := \{(\alpha, \beta) \in \Gamma''(m, r, s) : \#\beta = 1\}$
is equal to

$$\{(\underline{m}, \{i\}) : 1 \leq i \leq m\}$$

if $r = m$ and is equal to

$$\{(\underline{m} - \{i\}) : 1 \leq i \leq m\}$$

if $r = m - 1$ which it is empty otherwise.

Let $\Gamma^*(m, r, s) := \Gamma(m, r, s) - (\Gamma'(m, r, s) \cup \Gamma''(m, r, s))$.

Then $\Gamma^*(m, r, s)$ is empty, if $r + s < m$

and if $r + s \geq m$, then

$$\#\Gamma^*(m, r, s) = \binom{m}{r} \binom{r}{r+s-m} = \frac{m!}{(m-r)! (r+s-m)! (m-s)!}$$

Let $T, S, V \in \mathbb{P}$ with $ar(T) = m, ar(S) = r$

$ar(V) = s, T = T_1 \cdot \dots \cdot T_m, S = S_1 \cdot \dots \cdot S_r, V = V_1 \cdot \dots \cdot V_s, T_i, S_i, V_i \in \mathbb{P} - \{\underline{1}\}$ and

$\delta = (\alpha, \beta) \in \Gamma(m, r, s)$

Then

$\binom{T}{S, V} \gamma$ is defined to be the number of subsets I of $L(T) = L(T_1) \dot{\cup} \dots \dot{\cup} L(T_m)$
such that, if

$$\alpha := \{i \in \underline{m} : I \cap L(T_i) \neq \emptyset\}$$

$$\beta := \{i \in \underline{m} : I' \cap L(T_i) \neq \emptyset\}$$

where I' is the complement of I in $L(T)$ Then $\gamma = (\alpha, \beta) \in \Gamma(m, r, s)$ and
 $T \mid I = S, T \mid I' = V$.

Proposition 3.2. $\binom{T}{S, V} = \sum_{\gamma \in \Gamma(m, r, s)} \binom{T}{S, V} \gamma$

Proposition 3.3. (i) Let $\delta \in \Gamma'(m, r, s), \delta = (\{i\}, \underline{m} - \{i\})$. Then $s = m - 1$
and

$$\binom{T}{S, V} \delta = \binom{T_1}{\underline{1}, V_1} \cdot \dots \cdot \binom{T_{i-1}}{\underline{1}, V_{i-1}} \cdot \binom{T_i}{S, \underline{1}} \cdot \binom{T_{i+1}}{\underline{1}, V_i} \cdot \dots \cdot \binom{T_m}{\underline{1}, V_{m-1}}$$

(ii) Let $\delta \in \Gamma'(m, r, s), \delta = (\{i\}, \underline{m})$.

Then $s = m$ and

$$\binom{T}{S, V} \delta = \binom{T_1}{\underline{1}, V_1} \cdot \dots \cdot \binom{T_i}{S, V_i} \cdot \binom{T_{i+1}}{\underline{1}, V_{i+1}} \cdot \dots \cdot \binom{T_m}{\underline{1}, V_m}$$

- (iii) Let $\gamma \in \Gamma^*(m, r, s,), \gamma = (\alpha, \beta)$ Then $\#\alpha = r, \#\beta = s$ and $\binom{T}{S, V} \gamma = d_1 \cdot d_2 \cdot \dots \cdot d_m$
 where $d_i = \binom{T_i}{S_j, V_k}$ if $i \in \alpha \cap \beta$
 and $\alpha(j) = i, \beta(k) = i$ there $\alpha(j)$ is the j -th element of α and $\beta(k)$ is the k -th element of β in the natural order on $\underline{m}$.
 If $i \in \alpha, i \notin \beta$, then $d_i = \binom{T_i}{S_j, \underline{1}}$ with $\alpha(j) = i$
 If $i \in \beta, i \notin \alpha$, then $d_i = \binom{T_i}{\underline{1}, V_k}$ with $\beta(k) = i$.

4 Planar polynomials

Let K be a field and denote by $K\{x\}$ the K -algebra of planar polynomials. For any $T \in \mathbb{P}, T \neq \underline{1}, n = \deg(T)$, there is a K -multilinear operation $\cdot_T: (K\{x\})^n \rightarrow K\{x\}$ mapping $(S_1, \dots, S_n) \in \mathbb{P}^n$ onto the T -grafting $\cdot_T(S_1, \dots, S_n)$ which is a planar tree V in $\mathbb{P}$ containing T such that $V - \text{In}(T)$ is the planar forest $S_1 \dot{\cup} \dots \dot{\cup} S_n$ there $\text{In}(T)$ denotes the set of inner vertices of V .

Definition 4.1. Let $f \in K\{x\}$. Then

$$f^T: = \cdot_T(f, \dots, f)$$

is called the T -th power of F

Proposition 4.2. $(1 + x)^T = \sum_{S \in \mathbb{P}} \binom{T}{S} x^S$

Corollary 4.3. For any $f \in K\{x\}$ we have : $(1 + f)^T = \sum \binom{T}{S} f^S$

Denote by $\frac{d}{dx}: K\{x\} \rightarrow K\{x\}$ the derivation relative to x . It is a K -linear map satisfying the usual product rule for m -ary products such that $\frac{d}{dx}(x) = 1$

Proposition 4.4. For any $T \in \mathbb{P}, n = \deg(T)$,

$$\frac{d}{dx}(T(x)) = \sum_{\deg(U)=n-1} \binom{T}{U} x^U$$

Proposition 4.5. Let $T, S \in \mathbb{P}, T \neq 1 \neq S$.

Then for any $f \in K\{x\}$ we get

$$(f^T)^S = f^{(T^S)}$$

Corollary 4.6. $\left(\sum_{V \in \mathbb{P}} \binom{T}{V} x^V \right)^S = \sum \binom{T^S}{V} x^V$

5 Co-addition

Let $K\{x\} \otimes K\{x\}$ be the tensor product of the K -algebra $K\{x\}$ with itself. It is an algebra over the operad F freely generated by operation $(\mu_m)_{m \geq 2}$ where the degree of μ_m is m .

There is a unique K -algebra homomorphism

$$\Delta: K\{x\} \rightarrow K\{x\} \otimes K\{x\}$$

such that

$$\Delta(x) = x \otimes 1 + 1 \otimes x$$

Proposition 5.1. $\Delta(T(x)) = \sum_{S, V \in \mathbb{P}} \binom{T}{S} x^S \otimes x^V$

6 Planar roots

Let $K\{\{x\}\}$ be the K -algebra of planar power series in x .

Proposition 6.1. *Let $\text{char}(K) = 0$ and $T \in \mathbb{P}$.*

There is a unique power series

$$f \in K\{\{x\}\}$$

such that $f^T = (1+x)$ and whose constant term $f(0) = 1$.

We denote f by $\sqrt[T]{1+x}$ or $(1+x)^{T^{-1}}$ and call it the T -th root of $1+x$.

Example Let $T = x^2$ be the unique tree in $\mathbb{P}$ of degree 2. Then $\sqrt[T]{1+x}$ is denoted simply by $\sqrt{1+x}$.

One can show that $\sqrt{1+x} = \sum_{\substack{n=\deg(S)>0 \\ S \text{ binary}}} \frac{(-1)^{n-1}}{2^{2n-1}} x^S$

A tree S in $\mathbb{P}$ is binary, if the arity $\text{ar}_S(v)$ of each inner vertex of S is equal to 2. The arity $\text{ar}_S(v)$ is the number of edges of S which are incident with v and which are not on the path between v and the root of S .

Remark 6.2. One can show that

$$\sqrt[T]{1+x} = \exp_T\left(\frac{1}{n} \log_T(1+x)\right)$$

if $n = \deg(T)$ and $\exp_T$ is the exponential series relative to T which means that

$$(\exp_T(x))^T = \exp_T(nx)$$

$$\text{ord}(\exp_T(x) - 1 - x) \geq 2$$

and $\log_T$ is the inverse of $\exp_T$, see [G1], [G2] for the definitions of $\exp_T, \log_T$.

For any $T \in \mathbb{P}$, let $n_k(T)$ be the number of vertices of T of arity k . Then $n_0(T)$ is the degree of T and $n_1(T) = 0$ is equivalent to T being reduced.

If $T = T_1 \cdot \dots \cdot T_m, T_i \in \mathbb{P}, T_i \neq \underline{1}$ for all i , then

$$n_k(T) = \sum_{i=1}^m n_k(T_i)$$

if $k \neq m$ and

$$n_m(T) = 1 + \sum_{i=1}^m n_m(T_i).$$

One can show that

$$\sum_{k \geq 2} (k-1)n_k(T) = n_0(T) - 1$$

$$\text{Let } m(T) = \sum_{k=0}^{<00} n_k(T)$$

Proposition 6.3. *Let $C_d = x^d$ be the d -corona, $d \geq 2$, and*

$$\sqrt[d]{1+x}$$

the C_d -root of $(1+x)$.

Denote by $\binom{C_d^{-1}}{T}$ the coefficient of

$$\sqrt[d]{1+x}$$

relative to $T \in \mathbb{P}$.

Then $\binom{C_d^{-1}}{T} = 0$ if there is a vertex v in T with $ar_T(v) \geq d+1$.

If $ar_T(v) \leq d$ for all vertices v of T , then

$$\binom{C_d^{-1}}{T} = (-1)^{n(T)-n_0(T)} \cdot \prod_{k \geq 2} \left(\frac{1}{k} \binom{d-1}{k-1} \right)^{n_k(T)}$$

Let $v = (\nu_k)_{k \geq 2}$ be a sequence of integers $\nu_k \in \mathbb{N}$ such that $\nu_k = 0$ for almost all k .

Let $deg(\nu) = \nu_0 = 1 + \sum_{k=2}^{\infty} (k-1)\nu_k$ and $\mathbb{P}(\nu) = \{T \in \mathbb{P} : n_k(T) = \nu_k \text{ for all } k \geq 2\}$.

Then $\nu_0 = deg(T)$.

Denote by $C(\nu)$ the number of trees in $\mathbb{P}(\nu)$, it is called Catalan number of ν .

If $\nu_k = 0$ for $k \geq 3$, then $\mathbb{P}(\nu)$ is the set of binary trees of degree $\nu_0 = 1 + \nu_2$ and $C(\nu)$ is the usual Catalan number.

$$\text{Let } a_\nu = (-1)^{\nu-\nu_0} \prod_{k \geq 2} \left(\frac{1}{k} \binom{\nu_{0-1}}{k-1} \right)^{\nu_k}$$

Let $\binom{C_d^{-1}}{T} = a_{\nu(T)}$, if $\nu(T) = \left(n_k(T) \right)_{k \geq 2}$ by Proposition 6.3

Corollary 6.4. $\sum_{\deg(\nu)=n} \#C(\nu) \cdot a_\nu = \binom{1/d}{n} = (-1)^{n-1} \frac{1}{n!d^n} \prod_{r=0}^{n-1} (rd - 1)$

Proof. The canonical algebra homomorphism

$$\nu: K\{\{c\}\} \rightarrow K[[x]]$$

mapping T onto $x^{\deg(T)}$ maps $(1+x)^{c_d^{-1}}$ onto $(1+x)^{1/d} = \sum_{n=0}^{\infty} \binom{1/d}{n} x^n$. $\square$

Open problems: If the coefficient of $\sqrt[d]{1+x}$ relative to $S \in \mathbb{P}$ is denote by $\binom{\overline{T}^1}{S}$, then the question arises by what procedure this binomial coefficient $\binom{\overline{T}^1}{S}$ can be computed.

If there a recursive formula to compute $\binom{\overline{T}^1}{S}$.

More generally there is a power series f such that

$$f(0) = 1$$

$$f^T = (1+x)^S$$

One could denote f by $(1+S)^{S\overline{T}^1}$. How to compute the coefficients of f ?

References

- [G1] Gerritzen, L.: Planar rooted trees and non-associative exponential series, Advances in Applied Mathematics 33 (2004) 342-365
- [G2] Gerritzen, L.: Automorphisms of the planar tree power series algebra and the non-associative logarithm, Serdica Math. J. 30. Nos 2-3 (2004) 135-158
- [G3] Gerritzen, L.: Planar Shuffle Product, co-Addition and the non-associative Exponential, ArXiv:math.RA/0502378
- [G-H] Gerritzen, L. - Holtkamp, R.: Hopf co-addition for free magma algebras and the nonassociative Hausdorff series, J. of Algebra, 265 (2003), 264-284.
- [Hol] Holtkamp, R.: On Hopf algebra structures over operads, preprint, July 2004, ArXiv:math.RA/0407074.
- [R] Reutenauer, C.: Free Lie Algebras, Oxford Univ. Press, 1993
- [S] Stanley, R.: Enumerative Combinatorics I, II, Wadsworth-Brooks/Cole